

The cardinal characteristic for relative γ -setsArnold W. Miller ¹.

Abstract: For X a separable metric space define $\mathfrak{p}(X)$ to be the smallest cardinality of a subset Z of X which is not a relative γ -set in X , i.e., there exists an ω -cover of X with no γ -subcover of Z . We give a characterization of $\mathfrak{p}(2^\omega)$ and $\mathfrak{p}(\omega^\omega)$ in terms of definable free filters on ω which is related to the pseudointersection number $\mathfrak{p}$. We show that for every uncountable standard analytic space X that either $\mathfrak{p}(X) = \mathfrak{p}(2^\omega)$ or $\mathfrak{p}(X) = \mathfrak{p}(\omega^\omega)$. We show that both of following statements are each relatively consistent with ZFC: (a) $\mathfrak{p} = \mathfrak{p}(\omega^\omega) < \mathfrak{p}(2^\omega)$ and (b) $\mathfrak{p} < \mathfrak{p}(\omega^\omega) = \mathfrak{p}(2^\omega)$

First we define γ -set. An open cover $\mathcal{U}$ of a topological space X is an ω -cover iff for every finite $F \subseteq X$ there exists $U \in \mathcal{U}$ with $F \subseteq U$. The space X is a γ -set iff for every ω -cover of X there exists a sequence $(U_n \in \mathcal{U} : n < \omega)$ such that for every $x \in X$ for all but finitely many n we have $x \in U_n$, equivalently

$$X = \bigcup_{m < \omega} \bigcap_{n > m} U_n \quad \text{or} \quad \forall x \in X \forall^\infty n \in \omega \quad x \in U_n.$$

We refer to the sequence $(U_n : n < \omega)$ as a γ -cover of X , although technically we are supposed to assume that the U_n are distinct. In this paper all our spaces are separable metric spaces, so we may assume that all ω -covers are countable. This is because we can replace an arbitrary ω -cover with a refinement consisting of finite unions of basic open sets.

The γ -sets were first considered by Gerlits and Nagy [5]. One of the things that they showed was the following. The pseudointersection number $\mathfrak{p}$ is defined as follows:

$$\mathfrak{p} = \min\{|\mathcal{F}| : \mathcal{F} \subseteq [\omega]^\omega \text{ has the FIP and } \neg \exists X \in [\omega]^\omega \forall Y \in \mathcal{F} \ X \subseteq^* Y\}$$

¹Thanks for partial support from the BEST conference 2003, support from Boise State University, and to Tomek Bartoszyński, Justin Moore, and Marion Scheepers, for their hospitality during the time the main result in this paper was obtained.

Mathematics Subject Classification 2000: 03E35 54D20 03E50

Keywords: pseudointersection cardinal, relative γ -set, continuum, covering property, filter on ω

where FIP stands for the finite intersection property, i.e., every finite subset of $\mathcal{F}$ has infinite intersection, and $\subseteq^*$ denotes inclusion mod finite. The set X in this definition is called the pseudointersection of the family $\mathcal{F}$.

Gerlits and Nagy [5] showed that every γ -set has strong measure zero (in fact, the Rothberger property C''') and that Martin's Axiom implies every set of reals of size smaller than the continuum is a γ -set. Their arguments show that

$$\mathfrak{p} = \text{non}(\gamma\text{-set}) =^{def} \min\{|X| : X \text{ is not a } \gamma\text{-set}\}$$

where we only consider separable metric spaces X .

The property of being a γ -set is not hereditary. In fact, a γ -set X of size continuum is constructed in Galvin and Miller [4] using MA, which has the property that there exists a countable $F \subseteq X$ such that $X \setminus F$ is not a γ -set. However, any closed subspace of a γ -set is a γ -set.

Babinkostova, Guido and Kocinac [1] have defined the notion of a relative γ -set. This is also studied in Babinkostova, Kocinac, and Scheepers [2]. For $X \subseteq Y$ define X is a γ -set relative to Y iff for every open ω -cover $\mathcal{U}$ of Y there exists a sequence $(U_n \in \mathcal{U} : n < \omega)$ such that

$$X \subseteq \bigcup_{m < \omega} \bigcap_{n > m} U_n.$$

Note that if $Z \subseteq X \subseteq Y$ and X is a relative γ -set in Y , then Z is also.

Define the following cardinal number:

$$\mathfrak{p}(Y) = \min\{|X| : X \subseteq Y \text{ is not a } \gamma\text{-set relative to } Y\}.$$

Perhaps it should be written $\text{non}(\gamma \text{ relative to } Y)$.

In Just, Scheepers, Szeptycki, and Miller [8] many cardinal characteristics for covering properties are shown to be equal to well-known cardinals. Scheepers has noted that the cardinal numbers of the relativized version of the Rothberger property C''' work out to be either $\text{cov}(\text{meager})$ (the cardinality of the smallest cover of the real line with meager sets) or $\text{non}(\text{SMZ})$ (the cardinality of the smallest non strong measure zero set of reals).

Scheepers has raised the question of what we can say about the relativized versions for the γ -property. We begin with the easy

Proposition 1 $\mathfrak{p} \leq \mathfrak{p}(\omega^\omega) \leq \mathfrak{p}(2^\omega) \leq \mathfrak{c}$

Proof

If X is a γ -set, then it is a γ -set relative to any superspace. Let $|X| = \mathfrak{p}(\omega^\omega)$ be a subset of ω^ω which is not a relative γ -set. Then X is not a γ -set relative to itself, and hence $\mathfrak{p} \leq |X| = \mathfrak{p}(\omega^\omega)$.

For the second inequality, suppose $X \subseteq 2^\omega$ is not a γ -set relative to 2^ω with $|X| = \mathfrak{p}(2^\omega)$. Let $\mathcal{U}$ be an ω -cover of 2^ω witnessing that X is not a relative γ -set in 2^ω . Then

$$\{U \cup (\omega^\omega \setminus 2^\omega) : U \in \mathcal{U}\}$$

is an ω -cover of ω^ω witnessing that X is not a γ -set relative to ω^ω , and so $\mathfrak{p}(\omega^\omega) \leq |X| = \mathfrak{p}(2^\omega)$.

QED

We give another characterization of $\mathfrak{p}(\omega^\omega)$ and $\mathfrak{p}(2^\omega)$. A filter is free iff it contains the cofinite sets. For $\mathcal{F} \subseteq P(\omega)$ a free filter on ω , define

$$\mathfrak{p}_{\mathcal{F}} = \min\{|X| : X \subseteq \mathcal{F} \neg \exists a \in [\omega]^\omega \forall b \in X a \subseteq^* b\}.$$

Note that $\mathfrak{p}$ is the minimum of $\mathfrak{p}_{\mathcal{F}}$ for $\mathcal{F} \subseteq P(\omega)$ a free filter, since every family with the FIP generates a filter. We have the following characterizations:

Theorem 2 (a) $\mathfrak{p}(\omega^\omega)$ is the minimum of $\mathfrak{p}_{\mathcal{F}}$ such that $\mathcal{F} \subseteq P(\omega)$ is a Σ_1^1 free filter.

(b) $\mathfrak{p}(2^\omega)$ is the minimum of $\mathfrak{p}_{\mathcal{F}}$ such that $\mathcal{F} \subseteq P(\omega)$ is a Σ_2^0 free filter.

Proof

Suppose $X \subseteq \omega^\omega$ with $|X| = \mathfrak{p}(\omega^\omega)$ and $\mathcal{U}$ is an open ω -cover of ω^ω witnessing that X is not a relative γ -set. Without loss of generality we may assume that $\mathcal{U}$ is a countable family of clopen sets, say $\mathcal{U} = \{U_n : n \in \omega\}$. Let $f : \omega^\omega \rightarrow P(\omega)$ be the Marczewski [12] characteristic function of sequence

$$f(x) = \{n : x \in U_n\}.$$

This is a continuous mapping so its image is Σ_1^1 . Since $\mathcal{U}$ was an ω -cover the image has the FIP and note that the filter $\mathcal{F}$ generated by a Σ_1^1 family $\mathcal{G}$ with the FIP is Σ_1^1 , i.e.,

$$X \in \mathcal{F} \text{ iff } \exists F \in [\mathcal{G}]^{<\omega} \cap F \subseteq X.$$

Now assume $|X| < \mathfrak{p}(\mathcal{F})$. Then there exists $a \in \omega^\omega$ such that for each $b \in X$ we have that $a \subseteq^* f(b)$. It follows that $\langle U_n : n \in a \rangle$ is a γ -cover of X which is a contradiction. Hence $\mathfrak{p}(\omega^\omega) = |X| \geq \mathfrak{p}_{\mathcal{F}}$ and so

$$\mathfrak{p}(\omega^\omega) \geq \min\{\mathfrak{p}_{\mathcal{F}} : \mathcal{F} \text{ is a } \Sigma_1^1 \text{ free filter}\}.$$

To see that the other inequality suppose $\mathcal{F}$ is a Σ_1^1 filter and $X \subseteq \mathcal{F}$ witnesses $|X| = \mathfrak{p}_{\mathcal{F}}$. Let $f : \omega^\omega \rightarrow \mathcal{F}$ be a continuous onto map. For each $n \in \omega$ define $U_n = f^{-1}(\{x \in \mathcal{F} : n \in x\})$. Define $\mathcal{U} = \{U_n : n \in \omega\}$. Then $\mathcal{U}$ is an ω -cover of ω^ω . Choose $Y \subseteq \omega^\omega$ with $f(Y) = X$ and $|Y| = |X|$. If Y is relative γ in ω^ω , then there exists $a \in [\omega]^\omega$ such that $\langle U_n : n \in a \rangle$ is a γ -cover of Y . For each $b \in X$ we have $c \in Y$ with $f(c) = b$. For each n if $c \in U_n$, then $f(c) \in f(U_n)$ and so $n \in b$. It follows that $a \subseteq^* c$ for all $c \in X$. Since we are assuming that there is no such a , we must have that Y is not a γ -set relative to ω^ω and therefore

$$\mathfrak{p}(\omega^\omega) \leq |Y| = |X| = \min\{\mathfrak{p}_{\mathcal{F}} : \mathcal{F} \text{ is a } \Sigma_1^1 \text{ free filter}\}.$$

The proof for $\mathfrak{p}(2^\omega)$ is similar. Suppose $X \subseteq 2^\omega$ and $\mathcal{U}$ is an countable clopen ω -cover of 2^ω . Let $f : 2^\omega \rightarrow P(\omega)$ be $f(x) = \{n : x \in U_n\}$. Then f is continuous and so its range is a compact subset $C \subseteq P(\omega)$ which has the FIP. For each $n < \omega$ let $h : C^n \rightarrow P(\omega)$ be defined by

$$h(X_1, \dots, X_n) = X_1 \cap X_2 \cap \dots \cap X_n.$$

Then since h is continuous its range C_n is compact. Also since the projection of the compact set

$$\{(x, y) : x \in C_n \text{ and } x \subseteq y\}$$

onto the second coordinate is compact, we see that the filter $\mathcal{F}$ generated by C is Σ_2^0 in $P(\omega)$. Hence, if $X \subseteq 2^\omega$ has $|X| < \mathfrak{p}_{\mathcal{F}}$, then there exists $a \in [\omega]^\omega$ with $a \subseteq^* f(x)$ for each $x \in X$ and there for $x \in U_n$ for all but finitely many $n \in a$.

Conversely, suppose that $\mathcal{F}$ is a Σ_2^0 free filter in $P(\omega)$. Then there exists a compact $C \subseteq \mathcal{F}$ such that for every $x \in \mathcal{F}$ there exists a $y \in C$ with $x =^* y$. To see this, suppose that $\mathcal{F} = \cup_{n < \omega} C_n$. For each $n < \omega$ let

$$C_n^* = \{x \subseteq \omega : n \subseteq x \text{ and } \exists y \in C_n \forall i \geq n (i \in y \text{ iff } i \in x)\}$$

then $C = \cup_{n < \omega} C_n^*$ does the trick. Now suppose $X \subseteq \mathcal{F}$ with $|X| < \mathfrak{p}(2^\omega)$. Let $f : 2^\omega \rightarrow C$ be continuous and onto and choose $Y \subseteq 2^\omega$ with $|Y| = |X|$ such that for each $x \in X$ there exists $y \in Y$ with $f(y) =^* x$. Let $U_n = f^{-1}(\{x \in C : n \in x\})$. Then $\mathcal{U} = \{U_n : n < \omega\}$ is an ω -cover of 2^ω and

so there exists $a \in [\omega]^\omega$ such that for every $y \in Y$ we have that $y \in U_n$ for all but finitely many $n \in a$ and hence for each $x \in X$ there is $y \in Y$ with $a \subseteq^* f(y) =^* x$.

QED

For another paper studying the connection between γ -sets and free filters, see LaFlamme and Scheepers [10].

Lemma 3 (a) Suppose that X is homeomorphic to a closed subspace of Y , then $\mathfrak{p}(Y) \leq \mathfrak{p}(X)$.

(b) Suppose that $f : X \rightarrow Y$ is continuous and onto, then $\mathfrak{p}(X) \leq \mathfrak{p}(Y)$.

Proof

(a) Suppose $Z \subseteq X$ with $|Z| = \mathfrak{p}(X)$ is not relatively γ in X and this is witnessed by a family of open sets of Y which is an ω -cover of X . Then

$$\{U \cup (Y \setminus X) : U \in \mathcal{U}\}$$

is an ω -cover of Y which shows that Z is not relatively γ in Y . Hence $\mathfrak{p}(Y) \leq |Z| = \mathfrak{p}(X)$.

(b) Suppose $Z \subseteq Y$ with $|Z| = \mathfrak{p}(Y)$ is not relatively γ in Y and this is witnessed by an ω -cover $\mathcal{U}$. Choose $W \subseteq X$ with $|W| = |Z|$ and $f(W) = Z$. Let $\mathcal{V} = \{f^{-1}(U) : U \in \mathcal{U}\}$. Since f is onto, $\mathcal{V}$ is an ω -cover of X . We claim that there is no sequence $(f^{-1}(U_n) : n < \omega, U_n \in \mathcal{U})$ such that for every $x \in W$ for all but finitely many n we have $x \in f^{-1}(U_n)$. This is because $x \in f^{-1}(U_n)$ implies $f(x) \in U_n$, but then $f(W) = Z$ would have the property that every $y \in Z$ is in all but finitely many U_n . It follows that $\mathfrak{p}(X) \leq |W| = |Z| = \mathfrak{p}(Y)$.

QED

Theorem 4 Suppose X is an uncountable Σ_1^1 set in a Polish space, i.e., a nontrivial standard analytic space, then

- (a) if X is not σ -compact, then $\mathfrak{p}(X) = \mathfrak{p}(\omega^\omega)$ and
- (b) if X is σ -compact, then $\mathfrak{p}(X) = \mathfrak{p}(2^\omega)$.

Proof

If X is , it contains a homeomorphic copy of 2^ω and is the continuous image of ω^ω . It follows from Lemma 3 that

$$\mathfrak{p}(\omega^\omega) \leq \mathfrak{p}(X) \leq \mathfrak{p}(2^\omega).$$

(a) If X is not σ -compact, then Hurewicz [6] (see Kechris [9] 21.18) proved that there exists a closed subspace of X which is homeomorphic to ω^ω . Hence by Lemma 3 we have $\mathfrak{p}(X) \leq \mathfrak{p}(\omega^\omega)$.

(b) First suppose that $X = \omega \times 2^\omega$. Let $Y \subseteq \omega \times 2^\omega$ be non relatively γ with $|Y| = \mathfrak{p}(\omega \times 2^\omega)$. Since it is zero dimensional we can assume that this is witnessed by a countable ω -cover of clopen sets $\mathcal{U} = \{C_n : n < \omega\}$. As in the proof of Theorem 2 we take $f : \omega \times 2^\omega \rightarrow P(\omega)$ defined by

$$f(x) = \{n < \omega : x \in C_n\}.$$

The function f is continuous since the C_n are clopen and its image $f(\omega \times 2^\omega)$ is a σ -compact family of sets with the finite intersection property. Then $f(\omega \times 2^\omega)$ generates a σ -compact filter $\mathcal{F}$ and $f(Y)$ is a subset of $\mathcal{F}$ without a pseudo-intersection. Hence $\mathfrak{p}_{\mathcal{F}} \leq |f(Y)| \leq |Y| = \mathfrak{p}(\omega \times 2^\omega)$ and so we have $\mathfrak{p}(2^\omega) \leq \mathfrak{p}(\omega \times 2^\omega)$.

Now suppose that X is any σ -compact metric space. Note that there is a continuous onto mapping $f : \omega \times 2^\omega \rightarrow X$ and so by Lemma 3 we have that

$$\mathfrak{p}(X) \geq \mathfrak{p}(\omega \times 2^\omega) = \mathfrak{p}(2^\omega).$$

QED

The main result of this paper is the following theorem:

Theorem 5 *Both of following statements are each relatively consistent with ZFC:*

- (a) $\mathfrak{p} = \mathfrak{p}(\omega^\omega) < \mathfrak{p}(2^\omega)$ and
- (b) $\mathfrak{p} < \mathfrak{p}(\omega^\omega) = \mathfrak{p}(2^\omega)$

Proof

Part(a). The forcing notion we use is the obvious one. The only difficulty of the proof is that the forcing does only what we want it to and not more.

Given an ω -cover $\mathcal{U}$ of 2^ω define the poset $\mathbb{P}(\mathcal{U})$ as follows:

1. $p \in \mathbb{P}(\mathcal{U})$ iff $p = (F, (U_n \in \mathcal{U} : n < N))$ where $N < \omega$ and $F \in [2^\omega]^{<\omega}$.
2. $p \leq q$ iff $F^p \supseteq F^q$, $N^p \geq N^q$, $U_n^p = U_n^q$ for each $n < N^q$, and $x \in U_n^p$ for each $x \in F^q$ and n with $N^q \leq n < N^p$.

This poset is the obvious one for generically creating a γ -subcover of $\mathcal{U}$ for the ground model elements of 2^ω .

Lemma 6 *The partial order $\mathbb{P}(\mathcal{U})$ is σ -centered. Furthermore, suppose G is $\mathbb{P}(\mathcal{U})$ -generic over V . Define $(U_n : n < \omega)$ by $U_n = U_n^p$ for any $p \in G$ with $N^p > n$. Then $\forall x \in V \cap 2^\omega \ \forall^\infty n \ x \in U_n$.*

Proof

σ -centered is clear, since if $(N_n^p : n < N^p) = (N_n^q : n < N^q)$ then the condition $(F^p \cup F^q, (N_n^p : n < N^p))$ extends both p and q . The fact that U_n is defined for every $n < \omega$ follows from $\mathcal{U}$ being an ω -cover and a density argument, i.e., given any p with $N_p \leq n$ extend it by adding U_k which cover F_p . That $(U_n : n < \omega)$ is a γ -cover of $2^\omega \cap V$ since given any $x \in 2^\omega$ in the ground model, the set

$$D = \{p \in \mathbb{P}(\mathcal{U}) : x \in F_p\}$$

is dense and if $x \in F_p$ for some $p \in G$ then $x \in U_n$ for every $n \geq N_p$.

QED

The model for $\mathfrak{p} = \mathfrak{p}(\omega^\omega) < \mathfrak{p}(2^\omega)$ is obtained by starting with a model of GCH and doing a finite support iteration of $\mathbb{P}(\mathcal{U}_\alpha)$ for $\alpha < \omega_2$ where at each stage in the iteration

$$V[G_\alpha] \models \mathcal{U}_\alpha \text{ is an } \omega\text{-cover of } 2^\omega$$

and where we have dovetailed so as to ensure that for any $\mathcal{U}$ such that

$$V[G_{\omega_1}] \models \mathcal{U} \text{ is a countable } \omega\text{-cover of } 2^\omega$$

then for some $\alpha < \omega_2$ we have that $\mathcal{U} = \mathcal{U}_\alpha$. This dovetailing can be done since there are only continuum many countable ω -covers of 2^ω and the intermediate models satisfy the continuum hypothesis. In the model $V[G_{\omega_2}]$ we have that $\mathfrak{p}(2^\omega) = \omega_2$, so we need only show that $\mathfrak{p}(\omega^\omega) = \omega_1$. As usual, define Rothberger's unbounded number:

$$\mathfrak{b} = \min\{|X| : X \subseteq \omega^\omega \ \forall g \in \omega^\omega \ \exists f \in X \ \exists^\infty n \ f(n) > g(n)\}.$$

Lemma 7 $\mathfrak{p}(\omega^\omega) \leq \mathfrak{b}$

Proof

Suppose $X \subseteq \omega^\omega$ and $|X| < \mathfrak{p}(\omega^\omega)$. We need to show that X is eventually dominated. Without loss of generality we may assume that the elements of X are increasing and X is infinite. For each $n < \omega$ let

$$\mathcal{U}_n = \{U_m^n : m < \omega\} \text{ where } U_m^n = \{f \in \omega^\omega : f(n) < m\}.$$

Each $\mathcal{U}_n$ is an ω -cover of ω^ω . There is a standard trick due to Gerlits and Nagy [5] for replacing a sequence of ω -covers by a single ω -cover. Let

$$\{x_n : n < \omega\} \subseteq X$$

be distinct and let

$$\mathcal{U} = \{U \setminus \{x_n\} : n < \omega, U \in \mathcal{U}_n\}.$$

Then $\mathcal{U}$ is an ω -cover of ω^ω , since given any finite set F then $x_n \notin F$ for large enough n and so $F \subseteq U \setminus \{x_n\}$ for some $U \in \mathcal{U}_n$.

Since X is a relative γ -set in ω^ω there exists a sequence from $\mathcal{U}$ which is a γ -cover of X . Now since we threw out x_n from each element $\mathcal{U}_n$ at most finitely many of the elements of this sequence can come from the same $\mathcal{U}_n$. By taking an infinite subsequence we may assume that $(U_{g(n)}^n : n \in A)$ is a γ -cover of X for some infinite $A \subseteq \omega$. It follows that for every $f \in X$ that

$$\forall^\infty n \in A \quad f(n) < g(n).$$

Since the $f \in X$ are increasing if we extend g to all of ω by letting $g(m) = g(n)$ where $n \in A$ is minimal so that $n \geq m$, then g eventually dominates every $f \in X$ on all of ω .

It follows that $|X| < \mathfrak{b}$. Since X was arbitrary we get that $\mathfrak{p}(\omega^\omega) \leq \mathfrak{b}$.

QED

Our goal is to show that $\mathfrak{b} = \omega_1$ holds in this model. For the next two lemmas we assume $\mathcal{U}$ is an ω -cover of 2^ω and the forcing is $\mathbb{P}(\mathcal{U})$.

Lemma 8 *Suppose we are given $(U_n \in \mathcal{U} : n < N)$, $k < \omega$, and a term τ such that $\Vdash \tau \in \omega$. Then there exists $K < \omega$ such that for every $p \in \mathbb{P}(\mathcal{U})$ with $|F^p| \leq k$ and $(U_n \in \mathcal{U} : n < N) = (U_n^p \in \mathcal{U} : n < N^p)$ there exists $q \leq p$ such that $q \Vdash \tau < K$.*

Proof

Call $q \in \mathbb{P}(\mathcal{U})$ good iff

1. $N^q \geq N$
2. $U_n = U_n^q$ for all $n < N$, and
3. q decides τ , i.e. for some m , $q \Vdash \tau = m$.

For good q define:

$$U_q = \{(x_1, \dots, x_k) \in (2^\omega)^k : \forall i \ (N \leq i < N_q) \rightarrow \{x_1, \dots, x_k\} \subseteq U_i^q\}.$$

Note that each U_q is an open subset of $(2^\omega)^k$. Also the family U_q for q good cover $(2^\omega)^k$. This is because given any $(x_1, \dots, x_k)$ there exist a condition $q \leq (\{x_1, \dots, x_k\}, (U_n : n < N))$ which decides τ and therefor is good. By compactness there exist finitely many good q , say Γ , such that $\{U_q : q \in \Gamma\}$ covers $(2^\omega)^k$.

Since each good q decides τ and Γ is finite, we can find K so that for each $q \in \Gamma$

$$q \Vdash \tau < K.$$

Note that for any p as in the Lemma, if $F^p \subseteq U_q$ then q and p are compatible since $(F^p \cup F^q, (N_n^q : n < N^q))$ extends both of them.

QED

It is not hard to see from this lemma that our forcing does not add a dominating sequence. In order to prove the full result we need to show this for the iteration. To do this we prove the following stronger, but more technical, property (see Bartoszynski and Judah [3] definition 6.4.4).

Lemma 9 *The poset $\mathbb{P}(\mathcal{U})$ is really $\sqsubseteq^{\text{bounded}}$ -good, i.e., for every name τ for an element of ω^ω there exists $g \in \omega^\omega$ such that for any $x \in \omega^\omega$ if there exists $p \in \mathbb{P}(\mathcal{U})$ such that $p \Vdash \forall^\infty n \ x(n) < \tau(n)$, then $\forall^\infty n \ x(n) < g(n)$.*

Proof

Let $k_n, (U_m^n : m < N_n)$ for $n < \omega$ list with infinitely many repetitions all pairs of $k < \omega$ and finite sequences from $\mathcal{U}$. Using Lemma 8 repeatedly we can construct $g \in \omega^\omega$ such that for every $l < \omega$:

for any $n < l$ and $p \in \mathbb{P}(\mathcal{U})$ with

$$|F^p| \leq k_n \text{ and } (U_m^n : m < N_n) = (U_m^p : m < N^p)$$

there exists $q \leq p$ such that $q \Vdash \tau(l) < g(l)$.

Now suppose $p \Vdash \forall^\infty n \ x(n) < \tau(n)$. By extending p (if necessary) we may assume there exists n_0 such that $p \Vdash \forall n > n_0 \ x(n) < \tau(n)$. By making n_0 larger (if necessary) we may assume that

$$|F^p| = k_{n_0} \text{ and } (U_m^{n_0} : m < N_{n_0}) = (U_m^p : m < N^p).$$

Claim $\forall n > n_0 \ x(n) < g(n)$.

proof: Suppose not and $x(l) \geq g(l)$ for some $l > n_0$. By our construction of g we have that there exists $q \leq p$ such that $q \Vdash \tau(l) < g(l)$. But this means that $q \Vdash \tau(l) < x(l)$ which contradicts the fact that $p \Vdash \forall n > n_0 \tau(n) > x(n)$. This proves the Claim and the Lemma.

QED

It follows (see Bartoszyński and Judah [3] Theorem 6.5.4) that the finite support iteration using $\mathbb{P}(\mathcal{U}_\alpha)$ at stage α does not add a dominating real and so over a ground model which satisfies CH we have that $V[G_{\omega_2}]$ satisfies that $\mathfrak{b} = \omega_1$ and hence $\mathfrak{p}(\omega^\omega) = \omega_1$ by Lemma 7. This proves Theorem 5 part (a), the consistency of $\mathfrak{p} = \mathfrak{p}(\omega^\omega) < \mathfrak{p}(2^\omega)$.

Part (b) (the consistency of $\mathfrak{p} < \mathfrak{p}(\omega^\omega) = \mathfrak{p}(2^\omega)$) is simpler. It is well known that $\mathfrak{p} > \omega_1$ implies that $2^{\omega_1} = 2^\omega$. For example, see Rothberger [14]. Now starting with a ground model V which satisfies $2^\omega = \omega_2$ and $2^{\omega_1} = \omega_3$, do a finite support iteration using $\mathbb{P}(\mathcal{U}_\alpha)$ at stage $\alpha < \omega_2$ where $\mathcal{U}_\alpha$ is an ω -cover of $V[G_\alpha] \cap \omega^\omega$. Dovetail so that $\mathcal{U}_\alpha$ for $\alpha < \omega_2$ lists all countable ω -covers of ω^ω in the final model $V[G_{\omega_2}]$. This can be done since in all these models the continuum is ω_2 . The analogue of Lemma 6 holds for ω^ω in place of 2^ω so in the final model we have that $\mathfrak{p}(\omega^\omega) = \omega_2$. Also we get $\mathfrak{p} = \omega_1$ since $2^{\omega_1} = \omega_3 > \omega_2 = 2^\omega$. This finishes the proof of Theorem 5.

QED

One obvious question is

Question 10 *Is it consistent to have $\mathfrak{p} < \mathfrak{p}(\omega^\omega) < \mathfrak{p}(2^\omega)$?*

Question 11 *(Scheepers) Are either $\mathfrak{p}(\omega^\omega)$ or $\mathfrak{p}(2^\omega)$ the same as some other well-known small cardinal? See Vaughan [15] for a plethora of such cardinals.*

In Laver's model [11] for the Borel conjecture, we have that $\mathfrak{b} = \mathfrak{d} = \omega_2$ and $\mathfrak{p}(2^\omega) = \mathfrak{p}(\omega^\omega) = \omega_1$. In Laver's model there is a set of reals of size ω_1 which does not have measure zero, i.e., $\text{non}(\text{measure}) = \omega_1$, Judah and Shelah [7], see also Judah and Bartoszyński [3] or Pawlikowski [13]. But it is easy to see that $\mathfrak{p}(2^\omega) \leq \text{non}(\text{measure})$, i.e., if $X \subseteq 2^\omega$ and $|X| < \mathfrak{p}(2^\omega)$ then X has measure zero. Let $\{x_n : n < \omega\} \subseteq X$ be distinct and look at

$$\mathcal{U} = \{C \subseteq 2^\omega : \exists n \ x_n \notin C \text{ is clopen and } \mu(C) < \frac{1}{2^n}\}.$$

This is an ω -cover of 2^ω and so there exists a sequence $C_n \in \mathcal{U}$ with $X \subseteq \bigcup_n \bigcap_{m>n} C_m$. For any n at most finitely many C_n have measure $> \frac{1}{2^n}$ which shows that X has measure zero.

It is also true that $\mathfrak{p}(2^\omega) \leq \text{non}(\text{SMZ})$, i.e., if $|X| < \mathfrak{p}(2^\omega)$ then X has strong measure zero. The result of Gerlits and Nagy [5], that γ -sets have the Rothberger property C'' , relativizes to show that if $X \subseteq 2^\omega$ and $|X| < \mathfrak{p}(2^\omega)$, then X has the relative Rothberger property and this implies that X has strong measure zero.

Question 12 *Suppose that $Y = \bigcup_{n<\omega} X_n$ is an increasing union where Y is a separable metric space. If each X_n is relatively γ in Y , is Y a γ -set? If not, suppose each X_n is a γ -set, then is Y a γ -set?*

References

- [1] L. Babinkostova, C. Guido and Lj.D.R. Kocinac, On relative gamma sets, East-West Journal of Mathematics 2 (2000), 195 – 199.
- [2] L. Babinkostova, Lj.D.R. Kocinac, and Marion Scheepers, Combinatorics of open covers (VIII)
- [3] Bartoszyński, Tomek; Judah, Haim; **Set theory. On the structure of the real line.** A K Peters, Ltd., Wellesley, MA, 1995. xii+546 pp.
- [4] Galvin, Fred; Miller, Arnold W. γ -sets and other singular sets of real numbers. Topology Appl. 17 (1984), no. 2, 145–155.
- [5] Gerlits, J.; Nagy, Zs.; Some properties of $C(X)$. I. Topology Appl. 14 (1982), no. 2, 151–161.
- [6] Hurewicz, W.; Relativ perfekte teile von punktmengen und mengen (A); Fund. Math. 12 (1928), 78-109.
- [7] Judah, Haim; Shelah, Saharon; The Kunen-Miller chart (Lebesgue measure, the Baire property, Laver reals and preservation theorems for forcing). J. Symbolic Logic 55 (1990), no. 3, 909–927.
- [8] Just, Winfried; Miller, Arnold W.; Scheepers, Marion; Szeptycki, Paul J.; The combinatorics of open covers. II. Topology Appl. 73 (1996), no. 3, 241–266.

- [9] Kechris, Alexander S.; Classical descriptive set theory. Graduate Texts in Mathematics, 156. Springer-Verlag, New York, 1995.
- [10] Laflamme, Claude; Scheepers, Marion; Combinatorial properties of filters and open covers for sets of real numbers. *J. Symbolic Logic* 64 (1999), no. 3, 1243–1260.
- [11] Laver, Richard; On the consistency of Borel’s conjecture. *Acta Math.* 137 (1976), no. 3-4, 151–169.
- [12] Marczewski, E (Szpilrajn); The characteristic function of a sequence of sets and some of its applications, *Fund. Math.* 31(1938), 207-233.
- [13] Pawlikowski, Janusz; Laver’s forcing and outer measure. *Set theory* (Boise, ID, 1992–1994), 71–76, *Contemp. Math.*, 192, Amer. Math. Soc., Providence, RI, 1996.
- [14] Rothberger, Fritz; On some problems of Hausdorff and of Sierpiński. *Fund. Math.* 35, (1948). 29–46.
- [15] Vaughan, Jerry E.; Small uncountable cardinals and topology. With an appendix by S. Shelah. *Open problems in topology*, 195–218, North-Holland, Amsterdam, 1990. 54-02

Arnold W. Miller
 miller@math.wisc.edu
<http://www.math.wisc.edu/~miller>
 University of Wisconsin-Madison
 Department of Mathematics, Van Vleck Hall
 480 Lincoln Drive
 Madison, Wisconsin 53706-1388

The appendix is not intended for final publication but for the electronic version only.

Appendix Scheepers Remarks

Def. $X \subseteq Y$ is C'' (Rothberger) in Y iff for every sequence $(\mathcal{U}_n : n < \omega)$ of open covers of Y there exists a cover $(U_n \in \mathcal{U}_n : n < \omega)$ of X .

Prop.

- (a) $\text{non}(C'' \text{ in } \omega^\omega) = \text{non}(C'') = \text{cov}(\text{meager}) = \text{non}(\text{SMZ in } \omega^\omega)$
- (b) $\text{non}(C'' \text{ in } 2^\omega) = \text{non}(\text{SMZ in } 2^\omega) = \text{non}(\text{SMZ in } \mathbb{R})$

Proof

Here we mean strong measure zero in the usual metric on the reals and for ω^ω or 2^ω the metric $d(x, y) = \frac{1}{n+1}$ where n is minimal such that $x(n) \neq y(n)$.

(a) Fremlin and Miller (1988) prove:

- $\text{non}(C'') = \text{cov}(\text{meager}) = \text{non}(\text{SMZ in } \omega^\omega)$
- $\text{non}(C'') \leq \text{non}(C'' \text{ in } \omega^\omega)$ since if X is not relatively C'' it is not C'' .
- $\text{non}(C'' \text{ in } \omega^\omega) \leq \text{non}(\text{SMZ in } \omega^\omega)$ since $C'' \subseteq \text{SMZ}$.

(b) Suppose $X \subseteq 2^\omega$ fails to be relatively C'' . Note that by compactness of 2^ω we may assume there is a sequence $(\mathcal{U}_n : n < \omega)$ of finite clopen covers of 2^ω for which there is no $(U_n \in \mathcal{U}_n : n < \omega)$ which covers X . Now choose $\epsilon_n > 0$ so that any interval $[s]$ with diameter less than ϵ_n is a subset of some U_n . For the converse, suppose X fails to have SMZ in 2^ω which is witnessed by $(\epsilon_n : n < \omega)$. Then the sequence $\mathcal{U}_n = \{C \subseteq 2^\omega : \text{diam}(C) < \epsilon_n\}$ witnesses that it is not C'' .

$\text{non}(\text{SMZ in } 2^\omega) = \text{non}(\text{SMZ in } [0, 1]) = \text{non}(\text{SMZ in } \mathbb{R})$ is easy to prove.

QED

These cardinals can be different, for example, in the iterated modified Silver reals model, see Miller (1981), $\text{cov}(\text{meager}) = \omega_1$ while $\text{non}(\text{SMZ in } 2^\omega) = \omega_2$.

Def. X has the Menger property M iff for every sequence $(\mathcal{U}_n : n < \omega)$ of open covers there exists $(\mathcal{V}_n \in [\mathcal{U}_n]^{<\omega} : n < \omega)$ such that $X \subseteq \cup_n \mathcal{V}_n$.

Def. X has the Hurewicz property H iff for every sequence $(\mathcal{U}_n : n < \omega)$ of open covers there exists $(\mathcal{V}_n \in [\mathcal{U}_n]^{<\omega} : n < \omega)$ such that $X \subseteq \cup_m \cap_{n>m} \mathcal{V}_n$.

Then it is known, see Miller and Fremlin (1988) that

$$\mathfrak{d} = \text{non}(M) \quad \mathfrak{b} = \text{non}(H).$$

Relativizing H or M to 2^ω doesn't work since 2^ω has property H and M. For ω^ω it is easy to see:

Prop.

$$(a) \text{non}(M) \leq \text{non}(M \text{ in } \omega^\omega) \leq \mathfrak{d} \leq \text{non}(M)$$

$$(b) \text{non}(H) \leq \text{non}(H \text{ in } \omega^\omega) \leq \mathfrak{b} \leq \text{non}(H)$$

Biblio

Miller, Arnold W.; Some properties of measure and category. Trans. Amer. Math. Soc. 266 (1981), no. 1, 93–114.

Miller, Arnold W.; Fremlin, David H.; On some properties of Hurewicz, Menger, and Rothberger. Fund. Math. 129 (1988), no. 1, 17–33.

TeXed on : February 1, 2008